

# Vibrational spectroscopy of the borate mineral tunellite SrB<sub>6</sub>O<sub>9</sub>(OH)<sub>2</sub>·3(H<sub>2</sub>O) – Implications for the molecular structure



Ray L. Frost<sup>a,\*</sup>, Andrés López<sup>a</sup>, Ricardo Scholz<sup>b</sup>, Yunfei Xi<sup>a</sup>

<sup>a</sup> School of Chemistry, Physics and Mechanical Engineering, Science and Engineering Faculty, Queensland University of Technology, GPO Box 2434, Brisbane Queensland 4001, Australia  
<sup>b</sup> Geology Department, School of Mines, Federal University of Ouro Preto, Campus Morro do Cruzeiro, Ouro Preto, MG 35,400-00, Brazil

## HIGHLIGHTS

- Tunellite is a borate mineral of formula SrB<sub>6</sub>O<sub>9</sub>(OH)<sub>2</sub>·3(H<sub>2</sub>O).
- The structure is based upon B<sub>2</sub>O<sub>3</sub> units.
- Raman and infrared data on this mineral have been collected.
- The molecular structure of tunellite has been assessed by using vibrational spectroscopy.

## ARTICLE INFO

### Article history:

Received 19 August 2013

Accepted 7 October 2013

Available online 24 October 2013

### Keywords:

Raman spectroscopy

Borate

Tunellite

Infrared

Evaporite

## ABSTRACT

Tunellite is a strontium borate mineral with formula: SrB<sub>6</sub>O<sub>9</sub>(OH)<sub>2</sub>·3(H<sub>2</sub>O) and occurs as colorless crystals in the monoclinic pyramidal crystal system. An intense Raman band at 994 cm<sup>-1</sup> was assigned to the BO stretching vibration of the B<sub>2</sub>O<sub>3</sub> units. Raman bands at 1043, 1063, 1082 and 1113 cm<sup>-1</sup> are attributed to the in-plane bending vibrations of trigonal boron. Sharp Raman bands observed at 464, 480, 523, 568 and 639 cm<sup>-1</sup> are simply defined as trigonal and tetrahedral borate bending modes. The Raman spectrum clearly shows intense Raman bands at 3567 and 3614 cm<sup>-1</sup>, attributed to OH units. The molecular structure of a natural tunellite has been assessed by using vibrational spectroscopy.

© 2013 Elsevier B.V. All rights reserved.

## 1. Introduction

The mineral tunellite [1,2] is hydrated hydroxy borate of strontium of formula SrB<sub>6</sub>O<sub>9</sub>(OH)<sub>2</sub>·3H<sub>2</sub>O. The mineral is a rare mineral formed in borate deposits. The mineral is named after George Gerard Tunell, Jr. (1900–1996), Professor of Geology, University of California, Los Angeles, California, USA. The mineral is known from a significant number of deposits worldwide [1,3–6].

Tunellite is one member of a series of borate compounds of the borate series M<sup>2+</sup>O<sub>3</sub>B<sub>2</sub>O<sub>3</sub>·xH<sub>2</sub>O [7]. Other borate minerals include veatchite (Sr), p-veatchite (Sr) with x = 2, nobleite (Ca), with x = 4, aksaite (Mg) and gowerite (Ca) with x = 5 [7]. Many synthetic borates with this fundamental formula have been forthcoming. The chemistry of tunellite including its thermal decomposition has been studied by Gulensoy and Teberdar [8]. The effect of heat treatment on the structural properties of the tunellite has been studied [9]. Hydrogen bonding in tunellite has been elucidated

[10]. Boron NMR spectroscopy has proven useful for the study of the polyanions in tunellite [11].

Tunellite is monoclinic with point group 2/m [2,7]. The cell characteristics are P2<sub>1</sub>/a with a = 14.415(3), b = 8.213(1), c = 9.951(2), β = 114.05(1)° and Z = 4. Tunellite occurs in prismatic crystals up to 10 cm long or compact fine-grained nodules. The crystal structure contains infinite sheets composed of polymerized borate polyanions, with Sr<sup>2+</sup> cations and water molecules filling available spaces in and near the sheets [7]. Adjacent sheets are held together solely by bonds to water molecules [7]. The polymerized borate polyanions are composed of individual borate groups, each group containing three B–O tetrahedra and three B–O triangles.

Vibrational spectroscopy has been applied to borate glasses [12–15]. There have been a number of studies of borate glasses doped with a wide range of radioactive atoms [16,17]. Borate glasses are used as a means of containment of radioactive materials. There have been a number of studies looking at the effect of radiation on borate glasses [18,19]. If there is to be an understanding of borate glasses and their role in radioactive element containment, then an understanding of the vibrational spectroscopy of borate minerals needs to be undertaken. The number of vibrational

\* Corresponding author. Tel.: +61 73138 2407; fax: +61 73138 1804.  
 E-mail address: [r.frost@qut.edu.au](mailto:r.frost@qut.edu.au) (R.L. Frost).

spectroscopic studies of borate minerals is quite few and far between [20–23]. The number of Raman studies of borate minerals is also very limited [24,25]. There have been a number of infrared studies of some natural borates [26–29]. Most of these references are not new and there have been no recent studies on the vibrational spectroscopy of natural borates. Ross in Farmer's treatise reported the infrared spectra of several borate minerals [30].

The use of infrared spectroscopy is limited by the spatial resolution of the technique which is around 25  $\mu\text{m}$ . In comparison, the spatial resolution using Raman spectroscopy is 1  $\mu\text{m}$ . Thus, when studying a mineral using spectroscopic techniques it is advantageous to use Raman spectroscopy. The selection of the target mineral is more easily made. With infrared spectroscopy, any impurities will be measured as well as the target mineral. Raman spectroscopy has proven most useful for the study of secondary minerals [25,31–36].

To the best of the authors' knowledge, there have been very few infrared spectroscopic studies of tunellite type structure [29,37] and no Raman studies of this natural mineral have been forthcoming. The objective of this paper is to report the vibrational spectroscopic study of a natural tunellite mineral and relate the spectra to the molecular chemistry and the crystal chemistry of this borate mineral. We have characterised tunellite using a combination of Raman and infrared spectroscopy.

## 2. Experimental

### 2.1. Mineral

Details of the mineral have been published (page 80, vol. 5) [38].

### 2.2. Raman spectroscopy

Crystals of tunellite were placed on a polished metal surface on the stage of an Olympus BHSM microscope, which is equipped with 10 $\times$ , 20 $\times$ , and 50 $\times$  objectives. The microscope is part of a Renishaw 1000 Raman microscope system, which also includes a monochromator, a filter system and a CCD detector (1024 pixels). The Raman spectra were excited by a Spectra-Physics model 127 He–Ne laser producing highly polarised light at 633 nm and collected at a nominal resolution of 2  $\text{cm}^{-1}$  and a precision of  $\pm 1 \text{ cm}^{-1}$  in the range between 200 and 4000  $\text{cm}^{-1}$ . Repeated acquisitions on the crystals using the highest magnification (50 $\times$ ) were accumulated to improve the signal to noise ratio of the spectra. The spectra were collected over night. Raman Spectra were calibrated using the 520.5  $\text{cm}^{-1}$  line of a silicon wafer. The Raman spectrum of at least 10 crystals was collected to ensure the consistency of the spectra.

### 2.3. Infrared spectroscopy

Infrared spectra were obtained using a Nicolet Nexus 870 FTIR spectrometer with a smart endurance single bounce diamond ATR cell. Spectra over the 4000–525  $\text{cm}^{-1}$  range were obtained by the co-addition of 128 scans with a resolution of 4  $\text{cm}^{-1}$  and a mirror velocity of 0.6329  $\text{cm/s}$ . Spectra were co-added to improve the signal to noise ratio. The infrared spectra are given in the [supplementary information](#).

Spectral manipulation such as baseline correction/adjustment and smoothing were performed using the Spectralcalc software package GRAMS (Galactic Industries Corporation, NH, USA). Band component analysis was undertaken using the Jandel 'Peakfit' software package that enabled the type of fitting function to be selected and allows specific parameters to be fixed or varied

accordingly. Band fitting was done using a Lorentzian–Gaussian cross-product function with the minimum number of component bands used for the fitting process. The Lorentzian–Gaussian ratio was maintained at values greater than 0.7 and fitting was undertaken until reproducible results were obtained with squared correlations of  $r^2$  greater than 0.995.

## 3. Results and discussion

The Raman spectrum of tunellite in the 4000–100  $\text{cm}^{-1}$  region is displayed in Fig. 1a. This figure reports the position of the Raman bands and their relative intensity. It is noted that there are parts of the spectrum where no intensity is observed. Therefore, the spectrum is subdivided into sections in subsequent figures so that more detailed assessment of the spectrum can be made. In a similar way, the infrared spectrum of tunellite in the 4000–500  $\text{cm}^{-1}$  region is reported in Fig. 1b. The spectrum is not shown below 500  $\text{cm}^{-1}$ . The reason for this is that we are using a reflectance technique and the ATR cell absorbs all incident radiation. In a similar fashion to the Raman spectrum, the infrared spectrum is divided into sections depending upon the types of vibrations being observed.

The Raman spectrum of tunellite in the 1200–700  $\text{cm}^{-1}$  spectral range is reported in Fig. 2a. The infrared spectrum of tunellite in the 1200–700  $\text{cm}^{-1}$  spectral range is reported in Fig. 2b. The Raman spectrum in this spectral region is dominated by a sharp intense band at 994  $\text{cm}^{-1}$ . Bands of significantly lesser intensity are observed at 1043, 1063, 1082 and 1113  $\text{cm}^{-1}$ . The Raman band at 994  $\text{cm}^{-1}$  is assigned to the BO stretching vibration of the  $\text{B}_6\text{O}_9$  units. The Raman bands at 1043, 1063, 1082 and 1113  $\text{cm}^{-1}$  are attributed to the BOH in-plane bending modes. Iliev et al. determined the Raman spectrum of a synthetic cobalt tunellite [39].




Fig. 1. (a) Raman spectrum of tunellite (upper spectrum) over the 4000–100  $\text{cm}^{-1}$  spectral range and (b) infrared spectrum of tunellite (lower spectrum) over the 4000–500  $\text{cm}^{-1}$  spectral range.




Fig. 2. (a) Raman spectrum of tunellite over the 1200–700  $\text{cm}^{-1}$  spectral range and (b) infrared spectrum of tunellite over the 1000–650  $\text{cm}^{-1}$  spectral range.




Fig. 4. (a) Raman spectrum of tunellite (upper spectrum) in the 3800–3100  $\text{cm}^{-1}$  spectral range and (b) infrared spectrum of tunellite (lower spectrum) in the 3800–2800  $\text{cm}^{-1}$  spectral range.




Fig. 3. (a) Raman spectrum of tunellite (upper spectrum) in the 700–400  $\text{cm}^{-1}$  spectral range and (b) infrared spectrum of tunellite (lower spectrum) in the 400–100  $\text{cm}^{-1}$  spectral range.




Fig. 5. Infrared spectrum of tunellite over the 1800–1500  $\text{cm}^{-1}$  spectral range.

The symmetry species of some vibrational modes were determined. Iliev et al. [40] used Raman imaging to show the ferroelectric properties of tunellite type compounds. These workers [40] showed that tunellites exhibit a sequence of transitions from the high temperature paraelectric cubic phase to ferroelectric orthorhombic, monoclinic, trigonal phases, and finally to a monoclinic phase at low temperatures where both ferroelectric and magnetic orders coexist. Kim and Somoano determined the improper ferroelectric transition using Raman spectroscopy [41].

In contrast, the infrared spectrum of tunellite (Fig. 2b) shows complexity with many overlapping bands, making the assignment of the infrared bands very difficult. The intense infrared band at

960  $\text{cm}^{-1}$  with shoulders at 929, 947 and 985  $\text{cm}^{-1}$  is assigned to the trigonal borate stretching modes. The overlapping infrared bands at 784, 809, 844, 863 and 877  $\text{cm}^{-1}$  are attributed to the infrared tetrahedral borate stretching and bending vibrations. The series of bands from 669 through to 737  $\text{cm}^{-1}$  are related to trigonal borate bending modes.

The Raman spectra in the 700–400  $\text{cm}^{-1}$  and in the 300–100  $\text{cm}^{-1}$  spectral range are illustrated in Fig. 3a and b. Sharp Raman bands are observed at 464, 480, 523, 568 and 639  $\text{cm}^{-1}$ . These bands are simply defined as trigonal and tetrahedral borate bending modes. A series of infrared bands at 669, 679, 699, 708, 728 and 737  $\text{cm}^{-1}$  (Fig. 2b) are due to these bending modes. Strong Raman bands are found in the far wavenumber region at 150, 332 and 371  $\text{cm}^{-1}$  (Fig. 3b). These bands may be simply described as lattice vibrations.

Raman spectrum of tunellite in the 3800–2700  $\text{cm}^{-1}$  spectral range and the infrared spectrum of tunellite in the 3700–2500  $\text{cm}^{-1}$  spectral range are reported in Fig. 4a and b. According to the commonly accepted formula of tunellite  $\text{SrB}_6\text{O}_9(\text{OH})_2 \cdot 3(\text{H}_2\text{O})$ , both water or OH groups are found in the formula of the mineral. The Raman spectrum clearly shows intense Raman bands at 3567 and 3614  $\text{cm}^{-1}$ . These bands are assigned to the stretching vibrations of the OH units. Two OH-stretching bands in the Raman spectra of the mineral, at 3405 and 3494  $\text{cm}^{-1}$ , respectively, are indicative of two distinct hydrogen positions in the structure. Other strong intensity bands at 3243, 3282, 3324, 3369, 3383 and 3430  $\text{cm}^{-1}$  are found. These broader bands are assigned to water stretching vibrations.

The infrared spectrum in the 1800–1300  $\text{cm}^{-1}$  region is reported in Fig. 5. An infrared band centred upon 1677  $\text{cm}^{-1}$  is observed which may be band component analysed into several components at 1631, 1660 and 1727  $\text{cm}^{-1}$ . These bands are attributed to water bending vibrations. The intensity as observed in the infrared spectrum is quite strong with a broad band at 1631  $\text{cm}^{-1}$  found.

The development of these type of compounds such as tunellite rests with their ferromagnetic and pyro-electric properties [42]. The development of piezoelectric properties is time continuous [43,44]. Also a hydroxy compound has been synthesised [45], although the vibrational spectroscopy of this compound was not reported. In the infrared spectrum (Fig. 2b) a series of low intensity bands are observed.

#### 4. Conclusions

There are many borate minerals which have yet to have their vibrational spectrum determined and the molecular structure assessed in terms of their vibrational spectrum. In this work we have measured the Raman and infrared spectrum of tunellite, a hydrated hydroxy borate mineral of calcium with formula:  $\text{SrB}_6\text{O}_9(\text{OH})_2 \cdot 3(\text{H}_2\text{O})$ . The importance of the mineral tunellite rests with the chemistry of the compound and the range of isomorphous substitutions which can be made in order to produce specially selected ferroelectric, pyroelectric and piezoelectric properties.

The Raman spectrum in this spectral region is dominated by a sharp intense band at 994  $\text{cm}^{-1}$ . Bands of significantly lesser intensity are observed at 1043, 1063, 1082 and 1113  $\text{cm}^{-1}$ . The Raman band at 994  $\text{cm}^{-1}$  is assigned to the BO stretching vibration of the  $\text{B}_6\text{O}_9$  units.

#### Acknowledgments

The financial and infra-structure support of the School of Chemistry, Physics and Mechanical Engineering, Science and

Engineering Faculty, is gratefully acknowledged. The Australian Research Council (ARC) is thanked for funding the instrumentation. R. Scholz thanks to FAPEMIG – Fundação de Amparo à Pesquisa do estado de Minas Gerais, (Grant No. CRA – APQ-03998-10).

#### Appendix A. Supplementary material

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.molstruc.2013.10.012>.

#### References

- [1] R.C. Erd, V. Morgan, J.R. Clark, U.S. Geological Survey Professional Paper No. 424-C, 1961, p. 294.
- [2] H. Gulensoy, T. Teberdar, Bulletin of the Mineral Research and Exploration Institute of Turkey No. 79, 1972, p. 30.
- [3] O. Baysal, Bulletin of the Mineral Research and Exploration Institute of Turkey No. 79, 1972, p. 22.
- [4] W. Berdesinski, H. Meixner, *Abhandlungen* 104 (1965) 93.
- [5] C. Helvacı, *Miner. Deposita* 19 (1984) 217.
- [6] C. Helvacı, R.N. Alonso, *Turkish J. Earth Sci.* 9 (2000) 1.
- [7] J.R. Clark, *Am. Mineral.* 49 (1964) 1548.
- [8] H. Gulensoy, T. Teberdar, Bulletin of the Mineral Research and Exploration Institute of Turkey No. 79, 1972, p. 50.
- [9] M.S. Yilmaz, S. Piskin, *J. Chem. Soc. Pak.* 34 (2012) 526.
- [10] P.C. Burns, F.C. Hawthorne, *Can. Mineral.* 32 (1994) 895.
- [11] J.D. Cuthbert, W.T. MacFarlane, H.E. Petch, *J. Chem. Phys.* 43 (1965) 173.
- [12] I. Ardelean, S. Cora, *J. Mater. Sci.: Mater. Electron.* 19 (2008) 584.
- [13] I. Ardelean, S. Cora, *J. Optoelectron. Adv. Mater.* 12 (2010) 239.
- [14] I. Ardelean, L. Griguta, *J. Optoelectron. Adv. Mater.* 9 (2007) 2068.
- [15] I. Ardelean, V. Timar, *J. Optoelectron. Adv. Mater.* 10 (2008) 246.
- [16] F.H. El Batal, M.A. Azooz, A.A. El-Khesheh, *Trans. Indian Ceram. Soc.* 68 (2009) 81.
- [17] F.H. ElBatal, Y.M. Hamdy, *Trans. Indian Ceram. Soc.* 67 (2008) 193.
- [18] C. Rajyasree, P.M. Vinaya Teja, K.V.R. Murthy, D. Krishna Rao, *Physica B: Condensed Matter (Amsterdam, Netherlands)* 406, 2011, p. 4366.
- [19] B. Sumalatha, I. Omkaram, T.R. Rao, C.L. Raju, *J. Non-Cryst. Solids* 357 (2011) 3143.
- [20] M. Mir, J. Janczak, Y.P. Mascarenhas, *J. Appl. Crystallogr.* 39 (2006) 42.
- [21] I. Mitov, Z. Cherkezova-Zheleva, V. Mitrov, *J. Balkan Tribol. Assoc.* 4 (1998) 191.
- [22] A.S. Povarennykh, E.I. Nefedov, *Geologicheskii Zhurnal (Russian Edition)* 31 (1971) 13.
- [23] V.S. Suknev, E.N. Diman, *Zh. Prikl. Spektrosk.* 10 (1969) 326.
- [24] V.F. Ross, J.O. Edwards, *Chem. Boron Compd.* (1967) 15.
- [25] R.L. Frost, *J. Raman Spectrosc.* 42 (2011) 540.
- [26] M.V. Akhmanova, *Zh. Strukt. Khim.* 3 (1962) 28.
- [27] D.A. Long, R.T. Bailey, *Spectrochim. Acta* 19 (1963) 1607.
- [28] A. Vasko, *J. Serb. Czech J. Phys.* 17 (1967) 1110.
- [29] C.E. Weir, *J. Res. National Bureau Standards, Section A: Phys. Chem.* 70 (1966) 153.
- [30] V.C. Farmer, *Mineralogical Society Monograph 4: The Infrared Spectra of Minerals*, 1974.
- [31] R.L. Frost, S. Bahfenne, *J. Raman Spectrosc.* 42 (2011) 219.
- [32] R.L. Frost, S. Bahfenne, J. Cejka, J. Sejkora, J. Plasil, S.J. Palmer, E.C. Keeffe, I. Nemeč, *J. Raman Spectrosc.* 42 (2011) 56.
- [33] R.L. Frost, J. Cejka, J. Sejkora, J. Plasil, B.J. Reddy, E.C. Keeffe, *Spectrochim. Acta, A* 78 (2011) 494.
- [34] R.L. Frost, S.J. Palmer, *Spectrochim. Acta, A* 78 (2011) 1250.
- [35] R.L. Frost, S.J. Palmer, H.J. Spratt, W.N. Martens, *J. Mol. Struct.* 988 (2011) 52.
- [36] R.L. Frost, J. Sejkora, J. Cejka, J. Plasil, S. Bahfenne, E.C. Keeffe, *J. Raman Spectrosc.* 42 (2011) 534.
- [37] C. Helvacı, *Mercian Geol.* 6 (1978) 257.
- [38] J.W. Anthony, R.A. Bideaux, K.W. Bladh, M.C. Nichols, *Silica, Silicates, Mineral data Publishing, Tucson, Arizona*, 1995.
- [39] M.N. Iliiev, V.G. Hadjiev, M.E. Mendoza, J. Pascual, *Phys. Rev. B* 76 (2007) 214112/1.
- [40] M.N. Iliiev, V.G. Hadjiev, J. Iniguez, J. Pascual, *Acta Phys. Polonica* 116 (2009) 19.
- [41] Q. Kim, R.B. Somoano, *Ferroelectrics* 36 (1981) 431.
- [42] A.G. Castellanos-Guzman, ISAF '94, in: *Proceedings of the IEEE International Symposium on Applications of Ferroelectrics*, 9th, University Park, Pa., August 7–10, 1994, p. 6.
- [43] O. Crottaz, J.P. Rivera, H. Schmid, *J. Korean Phys. Soc.* 32 (1998) S1261.
- [44] M. Delfino, G.M. Loiacono, P.S. Gentile, *Inorg. Chim. Acta* 43 (1980) 59.
- [45] U. Werthmann, H. Gies, J. Glinnemann, T. Hahn, *Zeit. Krist.* 215 (2000) 393.