

## First report of *Ehrlichia ewingii* detected by molecular investigation in dogs from Brazil

L. S. Oliveira<sup>1</sup>, K. A. Oliveira<sup>1</sup>, L. C. Mourão<sup>1</sup>, A. M. Pescatore<sup>2</sup>, M. R. Almeida<sup>3</sup>, L. G. Conceição<sup>4</sup>, M. A. M. Galvão<sup>5</sup> and C. Mafra<sup>1</sup>

<sup>1</sup>Laboratório de Bioquímica e Biologia Molecular de Agentes Infeciosos e Parasitários, Departamento de Bioquímica e Biologia Molecular, Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil,

<sup>2</sup>Federal University of Switzerland, Zurich, Switzerland, <sup>3</sup>Laboratório de Infectologia Molecular Animal, Departamento de Bioquímica e Biologia Molecular, Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil, <sup>4</sup>Hospital Veterinário, Departamento de Medicina Veterinária, Universidade Federal de Viçosa, Viçosa, Minas Gerais, Brazil and <sup>5</sup>Departamento de Nutrição e Clínica Social, Universidade Federal de Ouro Preto, Ouro Preto, Minas Gerais, Brazil

### INTRODUCTION

The genus *Ehrlichia* encompasses a group of obligate intracellular bacteria that infect phagocytes, are transmitted by ticks and are related to ehrlichiosis in a number of animal species such as dogs, equines, ruminants, felines and humans. *Ehrlichia ewingii* is the causal agent of canine (CGE) and human (HGE) granulocytic ehrlichiosis and is mainly transmitted by ticks of the species *Amblyomma americanum*. CGE causes moderately severe disease and polyarthritis in chronically infected dogs. It has been described in dogs from the United States and Cameroon. *E. ewingii* has not been reported to have been propagated in cell culture and is usually identified by molecular techniques. This work aimed to verify by molecular assay the incidence of *E. ewingii* in dogs from Vicosa, Minas Gerais state, Brazil.

### MATERIAL AND METHODS

In order to detect *E. ewingii* in a population of dogs from Brazil, 100 blood samples were collected from dogs undergoing consultations or examinations at the Veterinary Hospital of the Federal University of Vicosa, Brazil, from April to May 2007. DNA was extracted from white blood cells by using the Dneasy Tissue Kit (QIAGEN, Valencia, CA, USA) according to the manufacturer's recommendations. These samples were tested for nested-PCR with genus-specific primers ECC (5'-

AGAACGAACGCTGGCGGCAAGC-3') and ECB (5'-CGTAT-TACCGCGGCTGCTGGCA-3') [1] and species-specific primers for *E. ewingii* EE5 (5'-CAATTCCTTAAATAGTCTCTGACTAT-TAG-3') [2] and HE3 (5'-TATAGGTACCGTCATTATCTTCCC-TAT-3') [3]. The primary amplification used as template 4 µL of DNA from each sample, and the secondary amplification used as template 1.5 µL of the product from the first reaction. The following conditions were used: one cycle of 94°C for 4 min, 30 cycles of 94°C for 30 s, 60°C (for the primary amplification) or 58°C (for the secondary amplification) for 30 s, 72°C for 1 min, followed by a cycle of 72°C for 5 min. *E. ewingii* presence was confirmed by another reaction with PCR primers Dsb-321 (5'-TTGCAAAATGATGTCTGAAGAT-ATGAAACA-3') and Dsb-671-ew (5'-CAGCTCCACCAAT-GAATGTATTTCCTCAA-3') [4] with DNA extracted from the canine blood samples that were positive by the nested-PCR. The conditions for this reaction were: one cycle of 94°C for 4 min, 40 cycles of 94°C for 30 s, 48°C for 30 s, 72°C for 1 min, followed by one cycle of 72°C for 5 min. PCR products were visualised in 1.2% agarose gel stained with ethidium bromide.

### RESULTS AND DISCUSSION

Five dogs infected by *E. ewingii* were found by nested-PCR and the reaction with primers that amplify a fragment of the dsb gene confirmed the presence of *E. ewingii* in these five samples. Four animals showed anaemia and one of them also showed thrombocytopenia. One animal infected with *E. ewingii* showed no haematological abnormality.

Formerly, the only recognised *Ehrlichia* species in Brazil were *Ehrlichia canis* and *Ehrlichia chaffeensis*, which makes this the first study to provide molecular evidence for canine infection caused by *E. ewingii* in the country. Trapp *et al.* [5] evaluated the serum antibody prevalence for *E. canis* in a population of dogs at the Veterinary Hospital of the State University of Londrina (Brazil) and did

Corresponding author and reprint requests: C. Mafra, Departamento de Bioquímica e Biologia Molecular, Av. P.H. Rolfs, s/n, Edifício CCBII, sala 213, Vicosa, MG, 36570-000, Brazil

E-mail: mafra@ufv.br

No conflicts of interest declared.

not exclude the possibility of infection by *E. ewingii* because the clinical signs observed in some dogs were more compatible with ehrlichiosis caused by *E. ewingii*.

In North America, *E. ewingii* is mainly transmitted by *A. americanum* ticks. As this species is not common in South America, other species of these ectoparasites should be involved in the transmission of *E. ewingii* to dogs in Brazil. Murphy *et al.* [3] found *R. sanguineus*, *A. americanum* and *Dermacentor variabilis* infected with this bacterium, suggesting that these tick species could be potential vectors for *E. ewingii*. Because *R. sanguineus* is a very common tick in Brazil, it is possible that this species is involved in the transmission of *E. ewingii* to dogs in this country.

## CONCLUSION

These results suggest that *E. ewingii* may be circulating among dogs in the studied region and show the need for further research in order to provide a better understanding of *E. ewingii* ecology in Brazil. Such studies would allow a better management of epidemiological surveillance aimed at the prevention of new *E. ewingii* infections.

## ACKNOWLEDGEMENTS

The authors would like to thank CNPq for a doctoral grant to L. S. Oliveira and scientific initiation grant to L. C. Mourão, and to CAPES for the doctoral grant to K. A. Oliveira. We are grateful to Dr David H. Walker of The University of Texas Medical Branch for partial financial support given to this project.

## REFERENCES

1. Dawson JE, Stallknecht DE, Howerth EW *et al.* Susceptibility of white-tailed deer (*Odocoileus virginianus*) to infection with *Ehrlichia chaffeensis*, the etiologic agent of human ehrlichiosis. *J Clin Microbiol* 1994; **32**: 2725–2728.
2. Kocan AA, Levesque GC, Whitworth LC, Murphy GL, Ewing SA, Barker RW. Naturally occurring *Ehrlichia chaffeensis* infection in coyotes from Oklahoma. *Emerg Infect Dis* 2000; **6**: 477–480.
3. Murphy GL, Ewing SA, Whitworth LC, Fox JC, Kocan AA. A molecular and serologic survey of *Ehrlichia canis*, *E. chaffeensis* and *E. ewingii* in dogs and ticks from Oklahoma. *Vet Parasitol* 1998; **79**: 325–339.
4. Doyle CK, Labruna MB, Breitschwerdt EB *et al.* Detection of medically important *Ehrlichia* by quantitative multicolor TaqMan real-time polymerase chain reaction of the *dsb* gene. *J Mol Diagn* 2005; **7**: 504–510.
5. Trapp SM, Dagnone AS, Vidotto O, Freire RL, Amude AM, de Moraes ASA. Seroepidemiology of canine babesiosis and ehrlichiosis in a hospital population. *Vet Parasitol* 2006; **140**: 223–230.